



Mapping Your Clients to Coverage Continuity Support

Step 1: Identify all active clients ages 18-64 covered by Medicaid expansion.

Step 2: On a monthly basis, use data to sort this group into three buckets (definitions are provided as an appendix):



Step 3: Consider the specific support needs of each of these buckets:



Step 4: Consider which of your programs provide case management, psych rehab, or peer support. Sort each bucket by these programs or services to understand what percent of each bucket will have their redetermination support needs met through their existing treatment plan and care team.

Step 5: Identify the remainder of each bucket. For this remainder population, stratify clients based on the level of clinical risk and the likelihood that they will be able to maintain coverage with staff support.

High Risk / Low Likelihood of Coverage	High Risk / High Likelihood of Coverage
Low Risk / Low Likelihood of Coverage	Low Risk / High Likelihood of Coverage

Step 6: Develop coverage continuity pathways for each of the stratification categories based on your organization's internal resources, state resources, and partner relationships.

EXAMPLE

High Risk / Low Likelihood of Coverage: Absorb Costs - Assess ongoing care needs and payment options - If possible, absorb cost of uncompensated care	High Risk / High Likelihood of Coverage: Intensive Support Provide intensive support to meet community engagement requirements
Low Risk / Low Likelihood of Coverage: Leverage Partners - Assess ongoing care needs and readiness for discharge - Engage MCO or other care partner to provide support	Low Risk / High Likelihood of Coverage: Minimal Support - Assess ongoing care needs and readiness for discharge - Provide supports to meet community engagement requirements, if resources allow. If not, engage MCO or other care partner